Estuaries Vol. 15, No. 4, p. 573 December 1992

Authors Index, Volume 15

Able, K. W., 171
Ayvazian, S. G., 368

Babione, M., 443
Bell, J. D., 360
Bildstein, K. L., 147
Birdsong, R. S., 392
Blood, E., 147
Bokuniewicz, H. J., 458
Brawley, J., 443
Burger, J., 239

Cabelli, V. J., 1
Callaway, J. C., 218
Chambers, R. M., 53, 349
Clark, M. W., 465
Cooper, K., 239
Correll, D. L., 431
Costa, J., 443
Cowan, J. H., Jr., 392
Crossman, R., 257

Dauer, D. M., 384
D'Avanzo, C., 443
Day, J. W., Jr., 421
Deegan, L. A., 368
DeMeo-Andreson, B., 443

Finn, J. T., 368
Fisher, F. M., 246
Foreman, K., 443
Franz, D. R., 193
Frederick, P., 147

Gammill, S. P., 122
Gaston, G. R., 227
Geoghegan, P., 251
Germano, J. D., 1
Geyer, W. R., 97
Gochfeld, D. J., 239
Gochfeld, M., 239
Good, R. E., 130
Gould, E., 289
Groffman, P. M., 83

Hager, S. W., 40
Harvey, J. W., 349
Hersh, D., 443
Hopkinson, C. S., Jr., 549
Hosier, P. E., 122
Houde, E. D., 392
Howarth, R. W., 18
Hsieh, Y. P., 414

Humphries, P., 327
Humphries, R., 529
Hussey, B. H., 59
Hyndes, G. A., 327

Jaworski, N. A., 83
Jordan, T. E., 431
Josselyn, M.N., 218

Kain, T., 257
Kalke, R. D., 307
Keller, A. A., 83
Keppel, R. G., 251
Kuropat, C., 289

Lajtha, K., 443
LaMontagne, M., 443
Lapointe, B. E., 75, 465
Largier, J. L., 26
Lazzari, M. A., 282
Lehman, P. W., 335
Lipsky, D., 239
Littler, D. S., 75
Littler, M. M., 75
Liu, J. T., 109
Luckenbach, M. W., 186
Luedke, D., 289

McComb, A. J., 529
McGlathery, K. J., 18
McNeill, S. E., 360

Madden, C. J., 421
Magnien, R. E., 497
Maples, R. S., 428
Marcomini, A., 517
Marino, R., 18
Mateja, G. B., 392
Mattson, M. T., 251
Mercaldo-Allen, R., 289
Monbet, Y., 563
Montagna, P. A., 307
Monteleone, D. M., 230

Nienhuis, P. H., 538
Nixon, S. W., 270

Odum, W. E., 59, 349
Orson, R. A., 130
Orio, A. A., 517
Ortega, S., 158
Orth, R. J., 186

Pavoni, B., 517
Peckol, P., 443
Pereira, J. J., 289
Potter, I. C., 327
Prager, J. C., 83
Priest, J. S., 392

Radach, G., 477
Ranasinghe, J. A., 384
Rebelo, J. E., 403
Reichle, J. J., 251
Rey, J. R., 257
Rhoads, D. C., 1
Rodi, A. J., Jr., 384
Rountree, R. A., 171

Safina, C., 239
Saliva, J. E., 239
Schemel, L. E., 40
Sellner, K. G., 497
Sfriso, A., 517
Shaffer, J., 257
Sham, C.-H., 443
Sharp, W. C., 392
Shreffler, D. K., 204
Signell, R. P., 97
Simenstad, C. A., 204
Simpson, R. L., 130
Stahl, R., 257
Stanley, D. W., 270
Stevenson, D. K., 282
Stevenson, L. H., 428
Summers, R. M., 497
Sutherland, J. P., 158

Tanacredi, J. T., 193
Taylor, D. I., 68
Thom, R. M., 204
Tomasko, D. A., 214
Tuttle, J. H., 298

Ulanowicz, R. E., 298

Valente, R. M., 1
Valiela, I., 429, 443, 429
Vaughn, C. C., 246

Weller, D. E., 431
Williams, L. E., Jr., 428

Yang, C. H., 414

Ziskowski, J., 289

© 1992 Estuarine Research Federation

0160-8347/92/040573-1$01.50/0

Estuaries Vol. 15, No. 4, p. 574–577 December 1992

Subject Index, Volume 15

abiotic vectors, 147
aboveground biomass, 218
aboveground uptake, 549
abundance measures, 403
Acadian zoogeographic province, 368
Acanthaluteres spilomelanurus, 360
Adriatic Sea, 517
algal blooms, 465
alkali-labile phosphate, 289
alkaline phosphatase, 75
Alosa aestivalis, 171, 368
Alosa fallax, 403
Alosa mediocris, 171
Alosa pseudoharengus, 171, 368
Alosa sapidissima, 171, 368
Altamaha River, Georgia, 549
Aluterus spp., 171
Ammodytes americanus, 171, 230, 368
Ammodytes tobianus, 403
ammonia, 53, 349, 443, 465, 477
ammonium, 538, 563
Ampelisca, 193
Ampelisca abdita, 193, 307
amphipods, 193
Anadara ovalis, 384
Anadara transversa, 384
Anas crecca, 227
Anchoa hepsetus, 171, 230, 368
Anchoa mitchilli, 171, 230, 368
Anguilla anguilla, 171, 368, 403
animal waste, 83
anthropogenic nutrient inputs, 18
Apeltes quadracus, 171, 230, 368
Aphia minuta, 403
aquarium studies, 214
aquifer, 458
artificial hydrologic regimes, 53
artificial salt-marsh, 257
Asabellides oculata, 384
Asychis elongata, 384
Atherina boyeri, 403
Atherina presbyter, 403
Atlantic cutlassfish, 251
Atlantic herring, 282
atmospheric deposition, 83
atmospheric inputs, 549
Australian estuaries, 327

Bahamas, 75
Bairdiella chrysoura, 171
Barrier Lagoon, Virginia, 349
beam trawls, 360
Belize, 75
Belone belone, 403
belowground biomass, 218
belowground production, 414
benthic algal production, 218
benthic communities, 193
benthic enrichment, 1
benthic habitat quality, 1
Bermuda, 18, 75
biogeochemical experiments, 53
biological uptake, 465
biomass, 403
biotic vectors, 147
bivalves, 384
black drum, 392
Black Rock Harbor, Connecticut, 289
blue crab, 186
blue-green algae, 529
Boston Harbor, Massachusetts, 289
bottom-water hypoxia, 270
Box-Jenkins transfer function models, 335
Brachaluteres jacksoniensis, 360
Brachidontes exustus, 307
brackish marshes, 431
Brevoortia tyrannus, 171, 230, 368

Cache River, Illinois, 549
calcium, 147
Callianassa sp., 307
Callinectes sapidus, 171, 186
Callinectes similis, 171
Callionymus lyra, 403
Cancer irroratus, 171
Cape Charles Harbor, Virginia, 392
Capitella capitata, 307
Caranx hippos, 171, 368
Carassius carassius, 403
carbon, 75
carbon flow, 298
carbonate-rich sediments, 18
carbonate-rich waters, 75
Carcinus maenus, 171
Carex, 549
Cassidinidea lunifrons, 307
catchments, 529
Centropogon australis, 360
Centropristis striata, 171, 368
Cerianthus americanus, 384
Chaetodon ocellatus, 171
Chaetopterus variopedatus, 384
change-in-storage, 83
Cheilodactylus vestitus, 360
Chelon labrosus, 403
Chesapeake Bay, 59, 298, 384, 392, 497
Chilomycterus schoepfi, 171, 230
chinook salmon, 204
chironomid insects, 204
chlorophyll *a*, 40, 230, 335, 421, 443, 563
chum salmon, 204
Ciliata mustela, 403
Clostridium perfringens, 1
Clupea harengus, 171, 282, 368
Clupeidae, 368
Clymenella mucosa, 307
Clymenella torquata, 384
coastal development, 443
coastal eutrophication, 465
coastal lagoons, 538
coastal lagoon system, 403
coastal land use, 431
coastal waters, 282
coastal watersheds, 443
Cobitus taenia, 403
conductance, 335
conductivity, 421
Conger oceanicus, 171
Connecticut River, 26
Conwy Estuary, Wales, 26
cordgrass, 414
Core Sounds, North Carolina, 158
cormorant, 327
Corophium louisianum, 307
Cossura delta, 307
couplings, 443
Crangon septemspinosa, 171
Crassostrea virginica, 158, 298
Cristiceps australis, 360
cropland, 431
ctenophore, 298, 392
Cyclaspis varians, 307
Cyclopterus lumpus, 368
Cynoscion regalis, 171, 230
Cyperaceae, 130
Cyprinodon variegatus, 171, 368
Cyrtopleura costata, 384

Delaware River basin, 130
Deltentosteus quadrimaculatus, 403
denitrification, 18, 83, 443
detritus, 218
detritus-based food chain, 204
diatom blooms, 529
diatoms, 477, 529
Dicentrarchus labrax, 403
diet comparisons, 327
Diodon nichthermerus, 360
Diopatra cuprea, 307, 384
Diplodus annularis, 403
Diplodus sargus, 403
Diplodus vulgaris, 403
dispersal of marine fishes, 251
dispersion, 97
dispersion patterns, 246
dissolved inorganic nitrogen, 40, 75, 465
dissolved nutrients, 349
dissolved organic carbon, 414
dissolved organic nirogen, 465
dissolved organic phosphorus, 465
dissolved oxygen, 270, 384, 403, 443, 477
dissolved reactive phosphorus, 40
Dutch estuaries, 538

eastern oyster, 158
Echiichthys vipera, 403
Edotea montosa, 307
Elbe River, 477
Elops saurus, 368
Ems River, 477
Enchelyopus cimbrius, 368

© 1992 Estuarine Research Federation

0160-8347/92/040574-4$01.50/0

enclosure experiments, 392
Engraulis encrasicholus, 403
enrichment, 68
epibenthic meiofauna, 227
episodic weather events, 109
Eteone heteropoda, 307, 384
Etropus microstomus, 230
Eudocimus albus, 147
Eulerian currents, 109
euryhaline annelids, 384
eutrophication, 1, 298, 465, 497, 517, 529, 538
evapotranspiration, 59
experimental chamber, 53
export, 335

fertilizer, 83, 443
fish assemblages, 368
fish fauna, 403
fish kills, 257
fish larvae, 282
fish stocks, 298
fish weir, 171
flagellates, 477
flathead, 327
flood control structures, 130
floodplain swamps, 431
flood tidal delta, 122
Florida Keys, 75, 465
food chain, 327
foraging, 204
forest swamps, 147
Fourleague Bay, Louisiana, 421
freckled blenny, 251
freshwater discharge, 257
freshwater inflow, 307
freshwater wetlands, 549
Fundulus diaphanus, 368
Fundulus heteroclitis, 171, 230, 368
Fundulus majalis, 171, 368

Gaidropsarus mediterraneus, 403
Galveston Bay, Texas, 246
Gambusia affinis, 403
Gammarus mucronatus, 307
Gasterosteus aculeatus, 171, 230, 368, 403
Gasterosteus sp., 368
Gasterosteus wheatlandi, 368
gelatinous predators, 392
German Bight, 477
global carbon dioxide levels, 549
Glycinde solitaria, 307, 384
Gobionellus boleosoma, 171
Gobiosoma bosci, 171, 368
Gobiosoma sp., 230
Gobius ater, 403
Gobius niger, 403
Gobius paganellus, 403
Gobius strictus, 403
Great Bay, New Jersey, 171
Great South Bay, New York, 230, 458
green-winged teal, 227
groundwater, 465
groundwater seepage, 458
groundwater tables, 130
growth rates, 392
Guadalupe Estuary, Texas, 307
guaguanche, 251
Gulf of Maine, 282
Gulf of Mexico, 421
Gymnapistes marmoratus, 360
Gyptis vittata, 307

habitat, 443
habitat selection, 186
habitat use, 368
Halodule wrightii, 214
Harvey Estuary, Western Australia, 529
Hemiramphus brasiliensis, 368
Hippocampus erectus, 171, 230
Hippocampus ramulosus, 403
Hobsonia florida, 307
Hog Island, Virginia, 349
Hudson River Estuary, 251
hydraulic theory, 26
hydrological environment, 403
hydromedusa, 392
Hyperoplus lanceolatus, 403
hypoxia, 270, 384
Hypsoblennius ionthas, 251

ichthyofauna, 230, 403
Illex illecebrosus, 171
impoundment, 257
incident photosynthetically active radiation, 421
Indian River Lagoon, Florida, 109, 257
industrial effluents, 517
infaunal community structure, 384
insects, 204
International Congress of Ecology, 429

Jamaica, 75
Jamaica Bay, New York, 193
James River, 384
Juncus roemerianus, 414
juvenile salmon, 204
juvenile winter flounder, 193

Kariega Estuary, South Africa, 68

Labrus viridis, 403
Lagocephalus lagocephalus, 403
Lagoon of Venice, 517
Lagrangian motions, 97
larval survival, 289
larval transport, 282
late Holocene, 130
leaching, 549
Leiostomus xanthurus, 171
Leitoscoloplos fragilis, 384
Leucon sp., 307
Libinia dubia, 171
Libinia emarginata, 171
life-history groups, 368
Limanda ferruginea, 368
limiting nutrient, 465
Limulus polyphemus, 171
Liopsetti putnami, 368
litter input, 257
Little Egg Harbor, New Jersey, 171
little pied, 327
Littoraria irrorata, 246
Littoridina sphinctostoma, 307
liver lesions, 289
living root biomass, 414
Liza aurata, 403
Liza ramada, 403
Liza slaiens, 403
Loch Creran, Scotland, 26
Long Island Sound, 289
long-term data sets, 270, 282, 335
Lophius americanus, 230
Lucania parva, 171, 368
Lutjanus griseus, 171

Macoma balthica, 384
Macoma mitchelli, 307
macroalga, 18, 75
macroalgae, 443, 517
macroalgal biomass, 443
macrobenthos, 384
Macrobrachiun intermedium, 360
Macroclymene zonalis, 384
macrofauna, 360
macrofaunal populations, 307
macrophytes, 59
macrotidal estuaries, 563
Mad Estuary, California, 26
maldanid polychaetes, 384
management, 122, 130, 204, 218, 239
management initiatives, 1
manmade canal system, 465
mapping, 421
marine invertebrates, 186
marsh comparisons, 257
mass action model, 298
mass balances, 83
measuring roots, 414
Mediomastus ambiseta, 384
Mediomastus californiensis, 307
medusae, 298
megalopae, 186
Megalops atlantica, 239
meiofauna, 227
meiofaunal populations, 307
Membras martinica, 171
Menidia beryllina, 171, 368
Menidia menidia, 230, 368
Menticirrhus saxatilis, 171, 230
Mercenaria mercenaria, 384
Merluccius bilinearus, 171
mesocosm, 68, 392
mesohaline ecosystem, 298
mesozooplankton, 298
microcosm, 53, 349
Microgadus tomcod, 368
Microphiopholis atra, 384
microtidal estuaries, 563
microzooplankton, 392
Mnemiopsis leidyi, 392
model, 335
modeling, 549
model predictions, 298
Moncanthus hispidus, 171, 251
Monoculodes nyei, 307
Morone americana, 171, 368
mortality rates, 392
mudflats, 431
Mugil cephalus, 171, 368, 403
Mugil curema, 171, 368
Mulinia lateralis, 307, 384

municipal sewage, 517
Mustelus canis, 171
Mya arenaria, 384
Myoxocephalus aenaeus, 232, 368

Narragansett Bay, Rhode Island, 1
native cordgrass, 218
Neanthes succinea, 307
Nectocarsinus tuberculosus, 360
needlerush, 414
Nemopsis bachei, 392
Neoodax baltealus, 360
Nereis succinea, 384
Nesogobius pulchellus, 360
nesting, 147
New Haven Harbor, Connecticut, 289
nitrate, 68, 335, 443, 465, 477, 538, 563
nitrogen, 18, 40, 75, 83, 147, 431, 477, 549, 563
Nodularia, 529
North Inlet Estuary, South Carolina, 147
North Sea, 477
Notropis bifrenatus, 368
Notropis heterolepis, 368
Nueces Estuary, Texas, 307
numerical model, 109
nutrient availability, 75
nutrient cycles, 517
nutrient cycling, 349, 549
nutrient enrichment, 18, 443
nutrient flux, 68, 431
nutrient fluxes, 477
nutrient input(s), 465, 477
nutrient limitation, 18
nutrient loadings, 193, 497, 538
nutrient pools, 497
nutrient processing, 53
nutrient release, 349
nutrient retention, 549
nutrient transport, 147, 431
nutrients, 18, 40, 53, 68, 83, 147, 335, 349, 443, 529, 538
Nyssa, 549

oceanic upwelling, 68
Oedalechilus labeo, 403
Okefenokee Swamp, Florida, 549
Oncorhynchus keta, 204
Oncorhynchus tshawytscha, 204
Ophidion marginata, 230
opportunistic species, 384
Opsanus tau, 171, 368
organic enrichment, 1
organic matter, 257, 349
Orthopristis chrysoptera, 251
Osmerus mordax, 368
Ovalipes ocelatus, 171
overharvesting, 298
oxygen demand, 1
oxygen evolution, 18
oyster management, 298
oyster stock rehabilition, 298

Pagellus bogaraveo, 403
Pagurus longicarpus, 171
Palaemonetes vulgaris, 171
paleoecological development, 130
Palmiet Estuary, South Africa, 26
Pamlico River Estuary, North Carolina, 270
Pamlico Sound, North Carolina, 158
Parablennius gattorugine, 403
Parablennius sanguinolentus, 403
Paralichthys dentatus, 171, 368
Parandalia ocularis, 307
Paraprionospio pinnata, 307, 384
particulate materials, 497
particulate organic carbon, 298
pasture, 431
Patuxent Estuary, 497
pectinaria gouldii, 307
Peel-Harvey estuarine system Western Australia, 529
Peel Inlet, Western Australia, 529
pelagic microbes, 298
Penaeus aztecus, 171
Penicillus capitatus, 18
Peprilus triacanthus, 230
Petromyzon marinus, 403
Phalocrocorax carbo, 327
Phalocrocorax melanoleucos, 327
Phalocrocorax sulcirostris, 327
Phalocrocorax varius, 327
phenology, 218
Phillips Creek, Virginia, 349
Phoeroides maculatus, 230
Pholis gunnellus, 368
Phoronis architecta, 307
phosphate, 53, 335, 349, 443, 477
phosphorus, 18, 40, 75, 83, 147, 431, 465, 477, 529, 549
phosphorus-binding capacity, 529
photosynthetic response, 18
physico-chemical variables, 421
phytoplankton, 443, 497, 563
pigfish, 251
planehead filefish, 251
Platichthys flesus, 403
Platycephalus speculator, 327
Pleuronectes americanus, 193, 230, 289
Pogonias cromis, 392
Pollachius virens, 171, 368
pollutant loading, 517
pollution, 1, 40
Polydora caulleryi, 307
Polydora ligni, 384
Polydora socialis, 307
polyhaline subtidal marsh, 171
Pomatochistus minutus, 403
Pomatomus saltatrix, 171, 368
porewater, 18, 257, 349
Posidonia australis, 360
potassium, 147
Potomac Estuary, 497
Potomac River, 83
predatory vertebrates, 327
prey density, 392
primary production, 214, 218, 298, 335, 421
Princeton-Jefferson Marsh, New Jersey, 130
Prionotus carolinus, 368
Prionotus evolans, 171, 368
Prionotus sp., 230
Pristigenys alta, 251
Procambaridae, 147
productivity, 75, 549, 563
Pseudopleuronectes, 193, 230
Pseudopleuronectes americanus, 171, 289, 368
Puerto Rican estuaries, 239
Pungitius pungitius, 368
Puyallup River estuary, Washington, 204
Pyramidella sp., 307

quantifying living roots, 414
Quercus-Castanea, 130

Rachycentron canadum, 171
raft culture, 298
Raja eglanteria, 171
Rangia cuneata, 307, 384
Rappanhannock River, 384
recruitment, 158, 392
Redfield proportions, 497
redox-potential discontinuity, 1
REMOTS®, 1
resource monitoring, 282
restoration, 218
restored wetland, 204
Rhine River, 477, 538
rhizome growth, 214
Rhode River estuary, 431
Ria de Aveiro, Portugal, 403
riparian forests, 431
riparian wetlands, 549
river exports, 83
river flow, 40
Rockefeller State Wildlife Refuge, Louisiana, 227
root biomass, 414
root uptake, 549

Sacramento River, 40
Sacramento-San Joaquin Delta, California, 335
salinity, 158, 307, 403, 421
salinity dilution, 497
salt front position, 251
saltmarsh development, 122
salt-marsh periwinkle, 246
salt-marsh surface, 68
sampling, 360
sampling gear, 171
San Francisco Bay, California, 40, 218
Sardina pilchardus, 403
Sardinella aurita, 171
Scobinichthys granulatus, 360
Scomber scombrus, 230
Scomberomorus sp., 171
Scophthalmus aquosus, 171, 230, 368
Scophthalmus rhombus, 403
seagrass, 360
seagrass meadows, 18, 465
sea-level rise, 122, 130
seawater inflow plunges, 26
Sebastian Inlet, Florida, 109
secchi disc, 335
secondary production 193, 307
sediment-profile photography, 1
sediment supply, 122

sedimentation, 158
sediment resuspension, 563
sediments, 1, 18, 257, 349
seine methodology, 171
Seiont Estuary, North Wales, 26
Selene vomer, 171
septic systems, 443
sessile organisms, 158
shear dispersion, 97
Sheepscot River, Maine, 282
shoal grass, 214
shorebird, 218
Shoreham, New York, 289
shoreline features, 97
short bigeye, 251
shrimp, 327
silicates, 477
siliciclastic waters, 75
sinks, 83
smooth cordgrass, 218
snails, 246
soil waterlogging, 122
Solea lascaris, 403
Solea senegalensis, 403
Solea vulgaris, 403
soluble reactive phosphorus, 68, 75
Spartina alterniflora, 122, 218, 349, 414
Spartina foliosa, 218
Sparus aurata, 403
spawning season, 392
species richness, 403
spectral composition of light, 214
Sphoeroides maculatus, 171, 368
Sphyraena borealis, 171
Sphyraena guachancho, 251
Spondyliosoma cantharus, 403
Stenotomus chrysops, 171, 368
Stigmatopora argus, 360
Stokes residual currents, 109
storms, 109
stratification, 26, 270
Streblospio benedicti, 307, 384
Strongylura marina, 171, 230, 368
subaqueous discharges, 458
Suisun Bay, California, 40, 335
sulfide, 257
Sundays Estuary, South Africa, 26
suspended solids, 335
swimming velocities, 186
Sygnanthus fuscus, 230
Symphodus bailloni, 403
Symphodus melops, 403
Syngnathus abaster, 403
Syngnathus acus, 403
Syngnathus fuscus, 171, 368
Syngnathus typhle, 403
Synodus foetens, 171

Tagelus divisus, 384
Tagelus plebeius, 307
Taurulus bubalis, 403
Tautoga onitis, 171, 230, 368
Tautogolabrus adspersus, 230, 368
Taxodium, 549
teleost, 327
temperature, 421
Thalassia testudinum, 214
tidal currents, 97
tidal dispersion, 97
tidal energy, 563
tidal fluctuations, 53
tidal freshwater, 384
tidal freshwater marshes, 53, 59, 130
tidal intrusion fronts, 26
tidal marshes, 53, 59
tidal mixing, 26, 563
tidal random walk, 97
tidal residual currents, 109
tides, 26
Tilapia mossambica, 239
topographic features, 97
Topsail Sound, North Carolina, 122
total dissolved nitrogen, 18
total dissolved phosphorus, 18
total nitrogen, 83, 497, 538
total phosphorus, 497, 538
toxicology, 289
Trachinotus falcatus, 171
translocation, 549
transparency, 403
transport, 349
trawls, 360
Trichiurus lepturus, 251
Trigla lucerna, 403
Trinectes maculatus, 171, 230, 368
trophic relationships, 298
turbidity, 335
turbulence, 97
turtle grass, 214

Ulva rigida, 517
Upeneichthys porosus, 360
upland forest, 431
upwelling, 68
Urophycis regia, 171
Urophycis sp., 171, 230
Urophysis tenuis, 368

Vaca Key, Florida, 214
vegetated habitats, 186
vegetation zonation, 130
Veerse Meer, The Netherlands, 538
Virginian zoogeographic province, 368
vitellogenin, 289
volatilization, 83

Wadden Sea, 538
wading bird, 147, 218
Waquoit Bay, Massachusetts, 368, 443
Washington, D.C., 83
wastewater treatment, 1, 40
water budgets, 549
water column, 53, 68
water-column stratification, 270
water loss, 59
water management, 538
watershed(s), 83, 443, 465
Wells Harbor, Maine, 368
Weser River, 477
wetlands, 59
white ibis, 147
wind mixing, 26
wind stress, 270
winter flounder, 289
Woodbury Creek Marsh, New Jersey, 130
Woods Hole, Massachusetts, 75
wrack deposition, 218

Xenanthura brevitelson, 307
xenobiotics, 289

yolk-sac larvae, 392
York River, 384